

## t59\_relset\_2 (TMWNsTaKTdgKbTHjuk- bjjW9H9JJjrHkTkAX)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_relat\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k6\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. k2\_relat\_1 (k2\_zfmisc\_1 X0 X1) = k2\_zfmisc\_1 X1 X0 \quad (1)$$

Assume the following.

$$\forall X0. (v1\_relat\_1 X0) \Rightarrow (\forall X1. (v1\_relat\_1 X1) \Rightarrow (k2\_relat\_1 (k6\_subset\_1 X0 X1) = k6\_subset\_1 (k2\_relat\_1 X0) (k2\_relat\_1 X1))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. ((m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))) \Rightarrow ((r2\_relset\_1 X0 X1 X2 X3) \Leftrightarrow (X2 = X3)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k6\_subset\_1 X0 X1 = k4\_xboole\_0 X0 X1 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (k3\_relset\_1 X0 X1 X2 = k2\_relat\_1 X2) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. v1\_relat\_1 (k2\_zfmisc\_1 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (m1\_subset\_1 (k3\_subset\_1 X0 X1) (k1\_zfmisc\_1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(m1\_subset\_1 (k3\_relset\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X1 X0))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0))\Rightarrow(k3\_subset\_1 X0 X1 = k4\_xboole\_0 X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (10)$$

**Theorem 1**

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(r2\_relset\_1 X1 X0 (k3\_relset\_1 X0 X1 (k3\_subset\_1 (k2\_zfmisc\_1 X0 X1) X2)) (k3\_subset\_1 (k2\_zfmisc\_1 X1 X0) (k3\_relset\_1 X0 X1 X2)))$$