

t5_arithm
(TMZ87SGoDcPXAa8TER3UR76crgKbjJoVgCR)

October 27, 2020

Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_xcmplx_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 X0 k6_numbers = k6_numbers) \quad (1)$$

Assume the following.

$$m1_subset_1 k6_numbers k1_numbers \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (v1_xcmplx_0 (k5_xcmplx_0 X0)) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (k7_xcmplx_0 X0 X1 = k3_xcmplx_0 X0 (k5_xcmplx_0 X1))) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k3_xcmplx_0 X0 X1 = k3_xcmplx_0 X1 X0) \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xcmplx_0 X0) \quad (6)$$

Theorem 1 $\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k7_xcmplx_0 k6_numbers X0 = k6_numbers)$.