

t5_arytm_1
(TMbNN33BCfz96cxp8VXf6YK5AMwAfoi21gr)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_arytm_2 : \iota$ be given. Let $r1_arytm_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k11_arytm_3 : \iota$ be given. Let $k7_arytm_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 \\ X1 k2_arytm_2) \Rightarrow (\neg(r1_arytm_2 X0 X1) \wedge (\forall X2.(m1_subset_1 \\ X2 k2_arytm_2) \Rightarrow (k7_arytm_2 X0 X2 \neq X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 \\ X1 k2_arytm_2) \Rightarrow ((k7_arytm_2 X0 X1 = k11_arytm_3) \Rightarrow (X0 = k11_arytm_3))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 \\ X1 k2_arytm_2) \Rightarrow (((r1_arytm_2 X0 X1) \wedge (X1 = k11_arytm_3)) \Rightarrow (X0 = \\ k11_arytm_3))) \end{aligned}$$