

t5_bvfunc_5
(TMQ4911z7DJHzb1BheSEZf6CyF14g7D41Q2)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_bvfunc_1 : \iota \Rightarrow \iota$ be given. Let $k9_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow ((r2_funct_2 X0 k6_margrel1 \\ & X2 (k12_bvfunc_1 X0)) \Rightarrow (r2_funct_2 X0 k6_margrel1 (k9_bvfunc_1 \\ & X0 X1 X2) (k12_bvfunc_1 X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow ((r2_funct_2 X0 k6_margrel1 \\ & (k9_bvfunc_1 X0 X1 X2) (k12_bvfunc_1 X0)) \Leftrightarrow (r1_bvfunc_1 X0 X1 X2))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow ((r1_bvfunc_1 X0 X1 X2) \Rightarrow (r1_bvfunc_1 \\ & X0 (k1_bvfunc_1 X0 X2) (k1_bvfunc_1 X0 X1)))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_funct_1 X1)\wedge (v1_funct_2 X1 X0 k6_margrel1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow(k1_bvfunc_1 X0 (k1_bvfunc_1 X0 X1) = X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_funct_1 X1)\wedge (v1_funct_2 X1 X0 k6_margrel1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow((v1_funct_1 (k1_bvfunc_1 X0 X1))\wedge((v1_funct_2 (k1_bvfunc_1 X0 X1) X0 k6_margrel1)\wedge(m1_subset_1 (k1_bvfunc_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \quad (5)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge (v1_funct_2 X1 X0 k6_margrel1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge ((v1_funct_2 X2 X0 k6_margrel1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow((r2_funct_2 X0 k6_margrel1 (k1_bvfunc_1 X0 X1) (k12_bvfunc_1 X0))\Rightarrow(r2_funct_2 X0 k6_margrel1 (k9_bvfunc_1 X0 X1 X2) (k12_bvfunc_1 X0))))$$