

t5_comseq_1
(TMZxmtEvxdJY9UJt8bGVM51F7aZAGBgpnM2)

October 27, 2020

Let $v2_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k2_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k5_complex1 : \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (r1_tarski X0 X1) \Rightarrow ((X2 \in X0) \vee (r1_tarski X0 (k4_xboole_0 X1 (k1_tarski X2)))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (2)$$

Assume the following.

$$k5_complex1 = k1_xboole_0 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v5_relat_1 X1 X0)) \Rightarrow (k2_relset_1 X0 X1 = k10_xtuple_0 X1) \quad (4)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow ((v2_relat_1 X0) \Leftrightarrow (\neg k1_xboole_0 \in k10_xtuple_0 X0)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow ((v5_relat_1 X1 X0) \Leftrightarrow (r1_tarski (k10_xtuple_0 X1) X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (8)$$

Theorem 1

$$\forall X0.(((v2_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v1_funct_2 X0 k5_numbers k2_numbers)\wedge(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers))))))\Rightarrow(r1_tarski (k2_relset_1 k2_numbers X0) (k6_subset_1 k2_numbers (k1_tarski k5_complex1)))$$