

t5\_decomp\_1  
(TMZfzB9EexrWLcbZDRCXcrmiTQZUGRqdxD4)

October 27, 2020

Let  $v2\_pre\_topc : \iota \Rightarrow o$  be given. Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v3\_decomp\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_decomp\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_pre\_topc : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tops\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Rightarrow (k3\_xboole\_0 X0 X1 = X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. r1\_tarski (k3\_xboole\_0 X0 X1) X0 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0)) \Rightarrow (k9\_subset\_1 X0 X1 X2 = k3\_xboole\_0 X1 X2) \quad (3)$$

Assume the following.

$$\forall X0. (l1\_pre\_topc X0) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow (k4\_decomp\_1 X0 X1 = k9\_subset\_1 (u1\_struct\_0 X0) X1 (k2\_pre\_topc X0 (k1\_tops\_1 X0 (k2\_pre\_topc X0 X1))))) \quad (4)$$

Assume the following.

$$\forall X0. (l1\_pre\_topc X0) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow ((v3\_decomp\_1 X1 X0) \Leftrightarrow (r1\_tarski X1 (k2\_pre\_topc X0 (k1\_tops\_1 X0 (k2\_pre\_topc X0 X1))))) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0)) \Rightarrow (k9\_subset\_1 X0 X1 X2 = k9\_subset\_1 X0 X2 X1) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. k3\_xboole\_0 X0 X1 = k3\_xboole\_0 X1 X0 \quad (7)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((v2\_pre\_topc\ X0)\wedge(l1\_pre\_topc\ X0))\Rightarrow(\forall X1. \\ & (m1\_subset\_1\ X1\ (k1\_zfmisc\_1\ (u1\_struct\_0\ X0)))\Rightarrow((v3\_decomp\_1 \\ & X1\ X0)\Leftrightarrow(X1 = k4\_decomp\_1\ X0\ X1))) \end{aligned}$$