

t5_hermitan

(TMWKPFTgv9cAv4rzRGznhK7TebFcB8hXRKM)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_complfld : \iota$ be given. Let $k2_complfld : \iota \Rightarrow \iota$ be given. Let $k4_complex1 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k15_complex1 : \iota \Rightarrow \iota$ be given. Let $k14_complex1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow ((X0 = k15_complex1 X0) \Rightarrow (k4_complex1 X0 = k6_numbers)) \quad (1)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 k1_complfld)) \Rightarrow (k2_complfld X0 = k14_complex1 X0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k15_complex1 X0 = k14_complex1 X0) \quad (3)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 k1_complfld)) \Rightarrow (v1_xcmplx_0 X0) \quad (4)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 k1_complfld)) \Rightarrow ((X0 = k2_complfld X0) \Rightarrow (k4_complex1 X0 = k6_numbers))$$