

t5\_int\_2

(TMQa7QUxhytNSBDBLasm2mDwWojJws7gvz3)

October 27, 2020

Let  $v1\_int\_1 : \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k3\_int\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_int\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_int\_1 X0) \Rightarrow ((r1\_int\_1 k6\_numbers X0) \Leftrightarrow (X0 = k6\_numbers)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_int\_1 X0) \wedge (v1\_int\_1 X1)) \Rightarrow (v7\_ordinal1 (k3\_int\_2 X0 X1)) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_int\_1 X0) \Rightarrow (\forall X1.(v1\_int\_1 X1) \Rightarrow (\forall X2. \\ & (v7\_ordinal1 X2) \Rightarrow ((X2 = k3\_int\_2 X0 X1) \Leftrightarrow ((r1\_int\_1 X2 X0) \wedge ((r1\_int\_1 \\ & X2 X1) \wedge (\forall X3.(v1\_int\_1 X3) \Rightarrow (((r1\_int\_1 X3 X0) \wedge (r1\_int\_1 \\ & X3 X1)) \Rightarrow (r1\_int\_1 X3 X2)))))))) \quad (3) \end{aligned}$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (v1\_int\_1 X0) \quad (4)$$

**Theorem 1**

$$\forall X0.(v1\_int\_1 X0) \Rightarrow (\forall X1.(v1\_int\_1 X1) \Rightarrow (((X0 = k6\_numbers) \wedge (X1 = k6\_numbers)) \Leftrightarrow (k3\_int\_2 X0 X1 = k6\_numbers)))$$