

t5_interva1
(TMFDoyk1zzbbxjTqERWriXhawtfSwWrvcJn)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_interval : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow ((\neg r1_tarski X1 X2) \Rightarrow (k1_interval \\ & X0 X1 X2 = k1_xboole_0))) \end{aligned} \tag{1}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow ((k1_interval X0 X1 X2 \neq k1_xboole_0) \Rightarrow \\ & (r1_tarski X1 X2))) \end{aligned}$$