

t5_jordan1a
(TMdPer2y4kMnVuzvQjV1etC7PAFwEYkz7ea)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k17_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $k9_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k25_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $v2_compts_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k24_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k10_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k13_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k6_pscomp_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (u1_struct_0 (k15_euclid np_2)))))) \Rightarrow ((X0 \in k17_pscomp_1 X1) \Rightarrow \\ & ((k18_euclid X0 = k18_euclid (k25_pscomp_1 X1)) \wedge ((v2_compts_1 \\ & X1 (k15_euclid np_2)) \Rightarrow ((r1_xxreal_0 (k17_euclid (k25_pscomp_1 \\ & X1)) (k17_euclid X0)) \wedge (r1_xxreal_0 (k17_euclid X0) (k17_euclid \\ & (k24_pscomp_1 X1)))))))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\ & np_2)))) \Rightarrow ((k18_euclid (k10_pscomp_1 X0) = k18_euclid (k25_pscomp_1 \\ & X0)) \wedge ((k18_euclid (k10_pscomp_1 X0) = k18_euclid (k24_pscomp_1 \\ & X0)) \wedge ((k18_euclid (k25_pscomp_1 X0) = k18_euclid (k24_pscomp_1 \\ & X0)) \wedge ((k18_euclid (k25_pscomp_1 X0) = k18_euclid (k13_pscomp_1 \\ & X0)) \wedge (k18_euclid (k24_pscomp_1 X0) = k18_euclid (k13_pscomp_1 \\ & X0)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid \\ & (k19_euclid X0 X1) = X0) \wedge (k18_euclid (k19_euclid X0 X1) = X1))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\ & np_2)))) \Rightarrow (m1_subset_1 (k9_pscomp_1 X0) k1_numbers) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (m1_subset_1 (k6_pscomp_1 X0) k1_numbers) \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (k10_pscomp_1 X0 = k19_euclid (k6_pscomp_1 X0) (k9_pscomp_1 X0)) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (7)$$

Theorem 1

$$\forall X0.((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow ((X1 \in k17_pscomp_1 X0) \Rightarrow (k18_euclid X1 = k9_pscomp_1 X0))$$