

## t5\_jordan23

(TMGhHkQ92dMbbTGH9FVqm9UcqrD1Kq4KaJh)

October 27, 2020

Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v2\_jordan23 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_1 : \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k7\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k4\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_card\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v6\_membered : \iota \Rightarrow o$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $v1\_card\_1 : \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((\neg r1\_xxreal\_0 (k1\_nat\_1 X1 np\_1) X0) \Leftrightarrow (r1\_xxreal\_0 X0 X1))) \quad (1)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow (((r1\_xxreal\_0 np\_1 X0) \wedge (r1\_xxreal\_0 (k1\_nat\_1 X0 np\_1) (k3\_finseq\_1 X1))) \Rightarrow ((X0 \in k4\_finseq\_1 X1) \wedge (k1\_nat\_1 X0 np\_1 \in k4\_finseq\_1 X1))) \wedge (((X0 \in k4\_finseq\_1 X1) \wedge (k1\_nat\_1 X0 np\_1 \in k4\_finseq\_1 X1)) \Rightarrow ((r1\_xxreal\_0 np\_1 X0) \wedge (r1\_xxreal\_0 (k1\_nat\_1 X0 np\_1) (k3\_finseq\_1 X1)))))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (3)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k4\_finseq\_1 X0 = k9\_xtuple\_0 X0) \quad (5)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k3\_finseq\_1 X0 = k1\_card\_1 X0) \quad (6)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k5\_numbers) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow ((\neg r1\_xxreal\_0 (k3\_finseq\_1 X1) X0) \Rightarrow ((k2\_nat\_1 X0 np\_1 \in k4\_finseq\_1 X1) \wedge (k1\_funct\_1 X1 (k2\_nat\_1 X0 np\_1) \in k10\_xtuple\_0 X1)))) \quad (7)$$

Assume the following.

$$v6\_membered k4\_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.(v1\_finset\_1 X0) \Rightarrow ((v1\_finset\_1 (k1\_card\_1 X0)) \wedge (v1\_card\_1 (k1\_card\_1 X0))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0) \Rightarrow ((v1\_funct\_1 X1) \wedge ((v1\_finseq\_1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0) \Rightarrow ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1) \wedge ((v5\_relat\_1 X1 X0) \wedge (v1\_funct\_1 X1))) \Rightarrow (\forall X2.(X2 \in k9\_xtuple\_0 X1) \Rightarrow (k7\_partfun1 X0 X1 X2 = k1\_funct\_1 X1 X2)) \quad (12)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow ((v2\_jordan23 X0) \Leftrightarrow (\forall X1.(m1\_subset\_1 X1 k5\_numbers) \Rightarrow (\neg(r1\_xxreal\_0 np\_1 X1) \wedge ((\neg r1\_xxreal\_0 (k3\_finseq\_1 X0) X1) \wedge (k1\_funct\_1 X0 X1 = k1\_funct\_1 X0 (k2\_nat\_1 X1 np\_1))))))) \quad (13)$$

Assume the following.

$$\forall X0.((v3\_ordinal1 X0)\wedge(v1\_finset\_1 X0))\Rightarrow(v7\_ordinal1 X0) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0)\Rightarrow(v5\_relat\_1 X1 X0) \quad (15)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow \\ ((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finset\_1 X0))) \quad (16)$$

Assume the following.

$$\forall X0.(v1\_card\_1 X0)\Rightarrow(v3\_ordinal1 X0) \quad (17)$$

Assume the following.

$$\forall X0.(v6\_membered X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 X0)\Rightarrow \\ (v7\_ordinal1 X1)) \quad (18)$$

**Theorem 1**

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0)\Rightarrow((v2\_jordan23 X1)\Leftrightarrow \\ (\forall X2.(m1\_subset\_1 X2 k5\_numbers)\Rightarrow(\neg(r1\_xxreal\_0 np\_1 \\ X2)\wedge((\neg r1\_xxreal\_0 (k3\_finseq\_1 X1) X2)\wedge(k7\_partfun1 X0 X1 X2 = \\ k7\_partfun1 X0 X1 (k2\_nat\_1 X2 np\_1)))))))$$