

t5_matrixr1

(TMS7jGZw6dJLZAQbQ2A2ejg38ypKrgdUcdY)

October 27, 2020

Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k8_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_euclid : \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k9_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k45_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k4_euclid : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v2_finseq_1 : \iota \Rightarrow o$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow (m2_finseq_2 X1 X0 (k4_finseq_2 (k3_finseq_1 X1) X0)) \quad (1)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (m2_finseq_2 X1 k1_numbers (k4_finseq_2 X0 k1_numbers)) \Rightarrow (k9_rvsum_1 X0 X1 (k5_finseq_2 k1_numbers X0 k6_numbers) = X1)) \quad (2)$$

Assume the following.

$$m1_subset_1 k1_xboole_0 k4_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0)\Rightarrow(\forall X2.(m2_finseq_2 X2 X0 X1)\Leftrightarrow(m1_subset_1 X2 X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Leftrightarrow(m1_finseq_1 X1 X0) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1 X0)\wedge((m1_subset_1 X1 (k4_finseq_2 X0 k1_numbers))\wedge(m1_subset_1 X2 (k4_finseq_2 X0 k1_numbers))))\Rightarrow(k9_rvsum_1 X0 X1 X2 = k45_valued_1 X1 X2) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v3_valued_0 X0)\wedge(v1_finseq_1 X0))))\wedge((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge((v3_valued_0 X1)\wedge(v1_finseq_1 X1))))))\Rightarrow(k8_rvsum_1 X0 X1 = k45_valued_1 X0 X1) \quad (8)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (9)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (10)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(k5_euclid X0 = k4_euclid X0) \quad (11)$$

Assume the following.

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\exists X1.(m1_finseq_1 X1 X0)\wedge((v1_relat_1 X1)\wedge((v4_relat_1 X1 k5_numbers)\wedge((v5_relat_1 X1 X0)\wedge((v1_funct_1 X1)\wedge((\neg v1_xboole_0 X1)\wedge((v1_finset_1 X1)\wedge((v1_finseq_1 X1)\wedge(v2_finseq_1 X1)))))))))) \quad (12)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (13)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Rightarrow(m1_subset_1 X2 X0)) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (16)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge((v7_ordinal1 X1)\wedge(m1_subset_1 X2 X0)))\Rightarrow(m2_finseq_2 (k5_finseq_2 X0 X1 X2) X0 (k4_finseq_2 X1 X0)) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.(v7_ordinal1 X0)\Rightarrow(m1_finseq_2 (k4_finseq_2 X0 X1) X1) \quad (19)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow((v1_relat_1 (k4_euclid X0))\wedge((v1_funct_1 (k4_euclid X0))\wedge((v1_finseq_1 (k4_euclid X0))\wedge(v3_valued_0 (k4_euclid X0)))))) \quad (20)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow(m2_subset_1 (k3_finseq_1 X0) k1_numbers k5_numbers) \quad (21)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(k4_euclid X0 = k5_finseq_2 k1_numbers X0 k6_numbers) \quad (22)$$

Assume the following.

$$\forall X0.\forall X1.(v1_xboole_0 X0)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_xboole_0 X2)) \quad (23)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v5_relat_1 X0 k1_numbers))\Rightarrow((v1_relat_1 X0)\wedge(v3_valued_0 X0)) \quad (24)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (25)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (26)$$

Assume the following.

$$\forall X0.(v6_membered X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(v7_ordinal1 X1)) \quad (27)$$

Theorem 1

$$\forall X0.(m2_finseq_1 X0 k1_numbers)\Rightarrow(k8_rvsum_1 X0 (k5_euclid (k3_finseq_1 X0)) = X0)$$