

t5_pboole (TMT- FrGte9BaYggauBKLHKK4ZEKBYXtGnSRc)

October 27, 2020

Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_pboole : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarSKI : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarSKI : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k4_tarSKI : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (X1 \in X0) \Rightarrow (k1_funct_1 (k2_funcop_1 X0 X2) X1 = X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarSKI (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarSKI X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k2_funcop_1 X0 X1)) \wedge (v1_funct_1 (k2_funcop_1 X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0. k1_pboole X0 = k7_funcop_1 X0 k1_xboole_0 \quad (5)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. ((X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (k4_tarSKI X1 X2 \in X0))) \wedge ((\neg X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \quad (6)$$

Theorem 1 $\forall X0. \forall X1. k1_funct_1 (k1_pboole X1) X0 = k1_xboole_0.$