

t5_prob_1 (TMQn-
FvXjoVPBwQ1HgkM75961fHgnsSDUUsV)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v2_finsub_1 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X1) \wedge (v2_finsub_1 X1) \wedge ((v1_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0)))))) \Rightarrow (k1_xboole_0 \in X1) \quad (1)$$

Assume the following.

$$\forall X0. k4_xboole_0 X0 k1_xboole_0 = X0 \quad (2)$$

Assume the following.

$$\forall X0. m1_subset_1 k1_xboole_0 (k1_zfmisc_1 X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k3_subset_1 X0 X1 = k4_xboole_0 X0 X1) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow ((v1_prob_1 X1 X0) \Leftrightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow ((X2 \in X1) \Rightarrow (k3_subset_1 X0 X2 \in X1)))) \quad (5)$$

Theorem 1

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X1) \wedge (v2_finsub_1 X1) \wedge ((v1_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0)))))) \Rightarrow (X0 \in X1)$$