

## t5\_radix\_2

(TMGp9eJwNtUAbvYHGvamTKyu6mZtmWXkzF)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k2\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v7\_ordinal1 X2) \Rightarrow ((X0 \in k2\_finseq\_1 X1) \Rightarrow (k2\_xcmplx\_0 X0 X2 \in k2\_finseq\_1 \\ & (k2\_xcmplx\_0 X1 X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \tag{2}$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v7\_ordinal1 X0) \wedge (m1\_subset\_1 X1 k5\_numbers)) \Rightarrow \\ & (k1\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \end{aligned} \tag{4}$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \tag{5}$$

### Theorem 1

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (( \\ & X0 \in k2\_finseq\_1 X1) \Rightarrow (k1\_nat\_1 X0 np\_1 \in k2\_finseq\_1 (k1\_nat\_1 \\ & X1 np\_1)))) \end{aligned}$$