

t5_semi_af1 (TMVe-
jGn5LMwoir9QMpU7YRHFpUCvVSDCnjm)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v1_semi_af1 : \iota \Rightarrow o$ be given. Let $l1_analoaf : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r2_analoaf : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0. ((\neg v2_struct_0 X0) \wedge ((v1_semi_af1 X0) \wedge (l1_analoaf \\
 & \quad X0))) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\
 & \quad (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3. (m1_subset_1 X3 \\
 & \quad (u1_struct_0 X0)) \Rightarrow (\forall X4. (m1_subset_1 X4 (u1_struct_0 X0)) \Rightarrow \\
 & \quad ((r2_analoaf X0 X1 X2 X3 X4) \Rightarrow (r2_analoaf X0 X3 X4 X1 X2))))))
 \end{aligned} \tag{1}$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_semi_af1 X0) \wedge (l1_analoaf \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 \\ & (u1_struct_0 X0)) \Rightarrow (\forall X4.(m1_subset_1 X4 (u1_struct_0 X0)) \Rightarrow \\ & ((r2_analoaf X0 X1 X2 X3 X4) \Rightarrow (r2_analoaf X0 X1 X2 X4 X3)))))) \end{aligned}$$