

t5_sin_cos7

(TMXtLLJu1f4SR8jt2z6CEHH6ujxEixhuX3Z)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_square_1 : \iota \Rightarrow \iota$ be given. Let $k3_square_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_square_1 : \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k7_binop_2 : \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 X1) \wedge ((\neg v2_xxreal_0 X0) \wedge (\neg v3_xxreal_0 X1)))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (r1_xxreal_0 k6_numbers (k3_xcmplx_0 X0 X0)) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X1 (k4_xcmplx_0 X0)) \wedge (r1_xxreal_0 (k2_xcmplx_0 X1 X0) k6_numbers))) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\neg r1_xxreal_0 (k7_square_1 (k9_binop_2 (k3_square_1 X0) np_1)) k6_numbers) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(r1_xxreal_0 X0 X1) \wedge ((\neg v2_xxreal_0 X1) \wedge (v2_xxreal_0 X0)))) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge (r1_xxreal_0 (k2_xcmplx_0 X1 X0) X1))) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(r1_xxreal_0 k6_numbers X0) \wedge ((\neg r1_xxreal_0 X1 X0) \wedge (r1_xxreal_0 (k6_square_1 X1) (k6_square_1 X0))))) \quad (8)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((r1_xxreal_0 X0 k6_numbers) \Rightarrow (k6_square_1 (k3_square_1 X0) = k4_xcmplx_0 X0)) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 X0 (k4_xcmplx_0 np_1) = k4_xcmplx_0 X0) \quad (10)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (11)$$

Assume the following.

$$(m2_subset_1 np_0 k1_numbers k5_numbers) \wedge ((m1_subset_1 np_0 k5_numbers) \wedge (m1_subset_1 np_0 k1_numbers)) \quad (12)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (13)$$

Assume the following.

$$k4_xcmplx_0 (k4_xcmplx_0 np_1) = np_1 \quad (14)$$

Assume the following.

$$k3_xcmplx_0 np_0 np_0 = np_0 \quad (15)$$

Assume the following.

$$k2_xcmplx_0 np_0 np_1 = np_1 \quad (16)$$

Assume the following.

$$r1_xxreal_0 (k4_xcmplx_0 np_1) np_0 \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(k9_binop_2 X0 X1 = k2_xcmplx_0 X0 X1) \quad (18)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(k7_square_1 X0 = k6_square_1 X0) \quad (19)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(k7_binop_2 X0 = k4_xcmplx_0 X0) \quad (20)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (21)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(k7_binop_2 (k7_binop_2 X0) = X0) \quad (22)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(v1_xreal_0 (k2_xcmplx_0 X0 X1)) \quad (23)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow((v1_xcmplx_0 (k4_xcmplx_0 X0))\wedge(v1_xreal_0 (k4_xcmplx_0 X0))) \quad (24)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xreal_0 (k3_square_1 X0)) \quad (25)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(m1_subset_1 (k7_square_1 X0) k1_numbers) \quad (26)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(m1_subset_1 (k7_binop_2 X0) k1_numbers) \quad (27)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0)\Rightarrow(k3_square_1 X0 = k3_xcmplx_0 X0 X0) \quad (28)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(k9_binop_2 X0 X1 = k9_binop_2 X1 X0) \quad (29)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0) \wedge (v3_xxreal_0 X0)) \Rightarrow ((\neg v1_xboole_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (\neg v2_xxreal_0 X0))) \quad (30)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (31)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0) \wedge (v2_xxreal_0 X0)) \Rightarrow ((\neg v1_xboole_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (\neg v3_xxreal_0 X0))) \quad (32)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (33)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (34)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\neg r1_xxreal_0 (k9_binop_2 (k7_square_1 (k9_binop_2 (k3_square_1 X0) np_1)) X0) k6_numbers)$$