

t5_topalg_3
(TMK5jgx B9FeZX8HpCrVh98hXsxEjUzp3v96)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_tex_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(l1_pre_topc X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (u1_struct_0 (k1_pre_topc X0 X1) = X1)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k6_domain_1 X0 X1 = k1_tarski X1) \quad (2)$$

Assume the following.

$$\forall X0.\neg v1_xboole_0 (k1_tarski X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(((\neg v2_struct_0 X0) \wedge (l1_pre_topc X0)) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0)))))) \Rightarrow ((\neg v2_struct_0 (k1_pre_topc X0 X1)) \wedge (v1_pre_topc (k1_pre_topc X0 X1))) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((l1_pre_topc X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow ((v1_pre_topc (k1_pre_topc X0 X1)) \wedge (m1_pre_topc (k1_pre_topc X0 X1) X0)) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.((\neg v2_struct_0 \\ & X2) \wedge ((v1_pre_topc X2) \wedge (m1_pre_topc X2 X0))) \Rightarrow ((X2 = k1_tex_2 X0 \\ & X1) \Leftrightarrow (u1_struct_0 X2 = k6_domain_1 (u1_struct_0 X0) X1)))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 \\ & (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((X2 = k1_tarski X1) \Rightarrow (k1_tex_2 \\ & X0 X1 = k1_pre_topc X0 X2)))) \end{aligned}$$