

t60_card_3

(TMF7mgn185sPkXixg5BNQ4AYEWQYWV3AK6E)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_card_3 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. k1_funct_1 (k16_funcop_1 X0 X1) X0 = X1 \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k16_funcop_1 X0 X1)) \wedge (v1_funct_1 (k16_funcop_1 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (v1_funct_1 (k7_funcop_1 X0 X1)) \wedge ((v1_funct_2 (k7_funcop_1 X0 X1) X0 (k1_tarski X1)) \wedge (m1_subset_1 (k7_funcop_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 (k1_tarski X1)))))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.k16_funcop_1 X0 X1 = k7_funcop_1 (k1_tarski X0) X1 \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(X1 = \\ k8_card_3 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.((v1_relat_1 \\ X3) \wedge (v1_funct_1 X3)) \wedge ((X2 = X3) \wedge (r1_tarski (k9_xtuple_0 X3) \\ (k9_xtuple_0 X0)) \wedge (\forall X4.(X4 \in k9_xtuple_0 X3) \Rightarrow (k1_funct_1 \\ X3 X4 \in k1_funct_1 X0 X4))))))) \quad (8) \end{aligned}$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarski X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (9)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow (((X0 \in k9_xtuple_0 X2) \wedge (X1 \in k1_funct_1 X2 X0)) \Rightarrow (k16_funcop_1 X0 X1 \in k8_card_3 X2))$$