

t60\_flang\_1  
(TMGeabnFoiqSemofJTp7PsddX9sEzGZQY5F)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k3\_catalan2 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k8\_flang\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k7\_flang\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k3\_catalan2 \\ & X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k3\_catalan2 \\ & X0))) \Rightarrow (\forall X3. (v7\_ordinal1 X3) \Rightarrow ((r1\_tarski X1 (k8\_flang\_1 \\ & X0 X2)) \Rightarrow (r1\_tarski (k7\_flang\_1 X0 X1 X3) (k8\_flang\_1 X0 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k3\_catalan2 X0))) \Rightarrow ((X1 \in k8\_flang\_1 X0 X2) \Leftrightarrow (\exists X3. (v7\_ordinal1 \\ & X3) \wedge (X1 \in k7\_flang\_1 X0 X2 X3))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (3)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k3\_catalan2 \\ & X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k3\_catalan2 \\ & X0))) \Rightarrow ((r1\_tarski X1 (k8\_flang\_1 X0 X2)) \Rightarrow (r1\_tarski (k8\_flang\_1 \\ & X0 X1) (k8\_flang\_1 X0 X2)))) \end{aligned}$$