

t60_qc_lang3 (TMXCmJj- exPmg1PngWAiT6mbSTNQZxVqrzMj)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k5_qc_lang3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k14_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k13_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ & X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (k9_qc_lang1 X0)) \Rightarrow (k5_qc_lang3 \\ & X0 (k14_qc_lang1 X0 X1 X2) = k4_subset_1 (k5_qc_lang1 X0) (k5_qc_lang3 \\ & X0 X1) (k5_qc_lang3 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ & X0)) \Rightarrow (k5_qc_lang3 X0 (k13_qc_lang1 X0 X1) = k5_qc_lang3 X0 X1)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((m1_qc_lang1 X0) \wedge ((m1_subset_1 \\ & X1 (k9_qc_lang1 X0)) \wedge (m1_subset_1 X2 (k9_qc_lang1 X0)))) \Rightarrow (m1_subset_1 \\ & (k14_qc_lang1 X0 X1 X2) (k9_qc_lang1 X0)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((m1_qc_lang1 X0) \wedge (m1_subset_1 X1 (k9_qc_lang1 \\ & X0))) \Rightarrow (m1_subset_1 (k13_qc_lang1 X0 X1) (k9_qc_lang1 X0)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ & X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (k9_qc_lang1 X0)) \Rightarrow (k2_qc_lang2 \\ & X0 X1 X2 = k13_qc_lang1 X0 (k14_qc_lang1 X0 X1 (k13_qc_lang1 X0 X2)))))) \end{aligned} \quad (5)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k9_qc_lang1\ X0)) \Rightarrow (\forall X2.(m1_subset_1\ X2\ (k9_qc_lang1\ X0)) \Rightarrow (k5_qc_lang3\ X0\ (k2_qc_lang2\ X0\ X1\ X2) = k4_subset_1\ (k5_qc_lang1\ X0)\ (k5_qc_lang3\ X0\ X1)\ (k5_qc_lang3\ X0\ X2)))) \end{aligned}$$