

t60_rfunct_3

(TMUZ7DwyXFaEJT9WQdwss3UR31QXRxLL3sG)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r2_rfunct_3 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_rfunct_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k15_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_funct_1 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ k1_numbers k1_numbers)))) \Rightarrow (\forall X1.\forall X2.(m1_subset_1 \\ X2 k1_numbers) \Rightarrow ((r2_rfunct_3 X0 X1) \Rightarrow (r2_rfunct_3 (k18_rfunct_3 \\ k1_numbers (k15_valued_1 k1_numbers k1_numbers X0 X2)) X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ r2_relset_1 X0 k1_numbers (k15_valued_1 X0 k1_numbers X1 k6_numbers \\ X1))) \end{aligned} \quad (3)$$

Assume the following.

$$(m2_subset_1 np_0 k1_numbers k5_numbers) \wedge ((m1_subset_1 np_0 \\ k5_numbers) \wedge (m1_subset_1 np_0 k1_numbers)) \quad (4)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((r2_rset_1 X0 X1 X2 X3)\Leftrightarrow(X2 = X3)) \quad (6)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (7)$$

Assume the following.

$$v3_membered k1_numbers \quad (8)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v3_membered X1)\wedge(((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\wedge(v1_xreal_0 X3)))\Rightarrow((v1_funct_1 (k15_valued_1 X0 X1 X2 X3))\wedge(m1_subset_1 (k15_valued_1 X0 X1 X2 X3) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (11)$$

Theorem 1

$$\forall X0.((v1_funct_1 X0)\wedge(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers k1_numbers))))\Rightarrow(\forall X1.(r2_rfunct_3 X0 X1)\Rightarrow(r2_rfunct_3 (k18_rfunct_3 k1_numbers X0) X1))$$