

t60_xxreal_1 (TM-
LuY6T1FC94iWCrh2axb7M7KWuYcVxRGWs)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_tarski (k1_xxreal_1 X0 X1) (k3_xxreal_1 X2 X3))) \Rightarrow ((\\ & \neg r1_xxreal_0 X0 X2) \wedge (r1_xxreal_0 X1 X3)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow ((r1_tarski (\\ & k2_xxreal_1 X0 X1) (k1_xxreal_1 X2 X3)) \Rightarrow ((r1_xxreal_0 X1 X0) \vee (\\ & r1_xxreal_0 X2 X0) \wedge (r1_xxreal_0 X1 X3)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow ((r1_xxreal_0 \\ & X0 X1) \Rightarrow ((r1_xxreal_0 X3 X2) \vee (r1_tarski (k1_xxreal_1 X1 X2) (k2_xxreal_1 \\ & X0 X3)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_xxreal_0 X2 X3)) \Rightarrow (r1_tarski (k3_xxreal_1 X1 X2) (k1_xxreal_1 \\ & X0 X3)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (r1_tarski \\ & (k3_xxreal_1 X0 X1) (k1_xxreal_1 X0 X1))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((r1_tarSKI X0 X1)\wedge(r1_tarSKI X1 X2))\Rightarrow(r1_tarSKI X0 X2) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0)\wedge(v1_xxreal_0 X1))\Rightarrow(r1_xxreal_0 X0 X0) \quad (7)$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0)\Rightarrow(\forall X1.(v1_xxreal_0 X1)\Rightarrow(\forall X2.(v1_xxreal_0 X2)\Rightarrow(\forall X3.(v1_xxreal_0 X3)\Rightarrow((r1_tarSKI (k2_xxreal_1 X0 X1) (k3_xxreal_1 X2 X3))\Rightarrow((r1_xxreal_0 X1 X0)\vee(\neg r1_xxreal_0 X0 X2)\wedge(r1_xxreal_0 X1 X3)))))))$$