

t60_xxreal_2
(TMQLNPcMohNx2rXNJjNftfiuiu38mnYacVN)

October 27, 2020

Let $v2_membered : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xxreal_2 : \iota \Rightarrow \iota$ be given. Let $m2_xxreal_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v2_membered X0) \Rightarrow (\forall X1.(v2_membered X1) \Rightarrow ((r1_tarski X0 X1) \Rightarrow (\forall X2.(m2_xxreal_2 X2 X1) \Rightarrow (m2_xxreal_2 X2 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.(v2_membered X0) \Rightarrow (v1_xxreal_0 (k2_xxreal_2 X0)) \quad (2)$$

Assume the following.

$$\forall X0.(v2_membered X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((X1 = k2_xxreal_2 X0) \Leftrightarrow ((m2_xxreal_2 X1 X0) \wedge (\forall X2.(m2_xxreal_2 X2 X0) \Rightarrow (r1_xxreal_0 X2 X1)))))) \quad (3)$$

Theorem 1

$$\forall X0.(v2_membered X0) \Rightarrow (\forall X1.(v2_membered X1) \Rightarrow ((r1_tarski X0 X1) \Rightarrow (r1_xxreal_0 (k2_xxreal_2 X1) (k2_xxreal_2 X0))))$$