

t61_chord
(TMZ3HQtTgL9cJHgo6zus3Ys4yZNAzRZrxy)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_glib_000 : \iota \Rightarrow o$ be given. Let $v4_glib_000 : \iota \Rightarrow o$ be given. Let $v2_chord : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k6_glib_000 : \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $r1_chord : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge ((v1_glib_000 X0) \wedge (v4_glib_000 X0)))))) \Rightarrow \\ (\exists X1. (m1_subset_1 X1 (k6_glib_000 X0)) \wedge (k6_glib_000 X0 = \\ k6_domain_1 (k6_glib_000 X0) X1)) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k6_domain_1 X0 X1 = k1_tarski X1) \quad (3)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\neg v1_xboole_0 (\\ k6_glib_000 X0)) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow ((v2_chord X0) \Leftrightarrow \\ (\forall X1. (m1_subset_1 X1 (k6_glib_000 X0)) \Rightarrow (\forall X2. (m1_subset_1 \\ X2 (k6_glib_000 X0)) \Rightarrow ((X1 \neq X2) \Rightarrow (r1_chord X0 X1 X2)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarSKI X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (6)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow ((v4_glib_000 X0) \Rightarrow (v2_chord X0))$$