

t61_compos_1
(TMP2EBipRdk9RLesqBXqQ6aHA32AgH929p8)

October 27, 2020

Let $l1_compos_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_compos_1 : \iota \Rightarrow \iota$ be given. Let $k2_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $k11_compos_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_afinsq_1 : \iota \Rightarrow o$ be given. Let $v5_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. (l1_compos_1 X1) \Rightarrow (\forall X2. (m1_subset_1 \\ X2 (u1_compos_1 X1)) \Rightarrow ((X0 \in k2_afinsq_1 (k11_compos_1 X1 X2)) \Leftrightarrow \\ ((X0 = k6_numbers) \vee (X0 = np_1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. ((\neg v1_xboole_0 X0) \wedge ((m1_subset_1 \\ X1 X0) \wedge (m1_subset_1 X2 X0))) \Rightarrow (k7_domain_1 X0 X1 X2 = k2_tarski X1 \\ X2) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((l1_compos_1 X0) \wedge (m1_subset_1 X1 (u1_compos_1 \\ X0))) \Rightarrow (((\neg v1_xboole_0 (k11_compos_1 X0 X1)) \wedge ((v1_relat_1 (k11_compos_1 \\ X0 X1)) \wedge ((v4_relat_1 (k11_compos_1 X0 X1) k5_numbers) \wedge ((v5_relat_1 \\ (k11_compos_1 X0 X1) (u1_compos_1 X0)) \wedge ((v1_funct_1 (k11_compos_1 \\ X0 X1)) \wedge ((v1_finset_1 (k11_compos_1 X0 X1)) \wedge (v1_afinsq_1 (k11_compos_1 \\ X0 X1)))))))))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v5_ordinal1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0)))) \Rightarrow (m1_subset_1 (k2_afinsq_1 X0) (k1_zfmisc_1 k5_numbers)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2_tarski X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 = X0) \vee (X3 = X1))) \quad (6)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_afinsq_1 X0))))) \Rightarrow ((v1_relat_1 X0) \wedge ((v5_ordinal1 X0) \wedge (v1_funct_1 X0))) \quad (7)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow ((v1_xboole_0 X1) \wedge ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)))) \quad (8)$$

Theorem 1

$$\forall X0. (l1_compos_1 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_compos_1 X0)) \Rightarrow (k2_afinsq_1 (k11_compos_1 X0 X1) = k7_domain_1 k5_numbers k6_numbers np_1))$$