

t61_eqrel_1 (TMZQeqN- VjsnBh1UhGmLr4yUbtbY5m3PN23x)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k3_tarSKI : \iota \Rightarrow \iota$ be given. Let $r1_tarSKI : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_setfam_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarSKI X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow (k5_setfam_1 X0 X1 = k3_tarSKI X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow (m1_subset_1 (k5_setfam_1 X0 X1) (k1_zfmisc_1 X0)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarSKI X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (4)$$

Theorem 1

$$\forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X1)) \Rightarrow (m1_subset_1 (k3_tarSKI X2) (k1_zfmisc_1 X0))))$$