

t61_flang_1
(TMG9tgsEDo3SkNcnZuvi7LzxkLJ4418CwFz)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k3_catalan2 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k3_catalan2 \\ & X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k3_catalan2 \\ & X0))) \Rightarrow ((r1_tarski X1 (k8_flang_1 X0 X2)) \Rightarrow (r1_tarski (k8_flang_1 \\ & X0 X1) (k8_flang_1 X0 X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k3_catalan2 X0))) \Rightarrow (r1_tarski X1 (k8_flang_1 X0 X1)) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \tag{3}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k3_catalan2 \\ & X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k3_catalan2 \\ & X0))) \Rightarrow ((r1_tarski X1 X2) \Rightarrow (r1_tarski (k8_flang_1 X0 X1) (k8_flang_1 \\ & X0 X2)))) \end{aligned}$$