

t61_rfunct_1

(TMX9vxKUHHsgBCDUyyUtfzWSWaRXwG36ooM)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_rfunct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k4_funct_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. ((m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow ((r2_relset_1 X0 X1 X2 X3) \Leftrightarrow (X2 = X3)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k7_rfunct_1 X0 X1 = k4_funct_3 X0 X1 \quad (4)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(v1_funct_1 (k7_rfunct_1 X0 X1))\wedge(m1_subset_1 (k7_rfunct_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X1 k1_numbers))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X2)\wedge(v1_funct_1 X2))\Rightarrow((X2 = k4_funct_3 X0 X1)\Leftrightarrow((k9_xtuple_0 X2 = X1)\wedge(\forall X3.(X3 \in X1)\Rightarrow(((X3 \in X0)\Rightarrow(k1_funct_1 X2 X3 = np_1))\wedge((\neg X3 \in X0)\Rightarrow(k1_funct_1 X2 X3 = k1_xboole_0)))))) \quad (8)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.\forall X2.((X1 \in k9_xtuple_0 X0)\Rightarrow((X2 = k1_funct_1 X0 X1)\Leftrightarrow(k4_tarski X1 X2 \in X0)))\wedge((\neg X1 \in k9_xtuple_0 X0)\Rightarrow((X2 = k1_funct_1 X0 X1)\Leftrightarrow(X2 = k1_xboole_0)))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (11)$$

Theorem 1

$$\forall X0.\forall X1.(\neg v1_xboole_0 X1)\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k1_numbers))))\Rightarrow((r2_relset_1 X1 k1_numbers X2 (k7_rfunct_1 X0 X1))\Leftrightarrow((k1_relset_1 X1 X2 = X1)\wedge(\forall X3.(m1_subset_1 X3 X1)\Rightarrow(((X3 \in X0)\Rightarrow(k1_funct_1 X2 X3 = np_1))\wedge((\neg X3 \in X0)\Rightarrow(k1_funct_1 X2 X3 = k6_numbers)))))))$$