

t62_abc Miz_1
(TMRjZ2P9ELQ7QS6XU4etbiFpYySX3LsEmiE)

October 27, 2020

Let $v1_instalg1 : \iota \Rightarrow o$ be given. Let $v1_abc Miz_1 : \iota \Rightarrow o$ be given. Let $v3_abc Miz_1 : \iota \Rightarrow o$ be given. Let $l1_msualg_1 : \iota \Rightarrow o$ be given. Let $v6_abc Miz_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_abc Miz_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k13_abc Miz_1 : \iota \Rightarrow \iota$ be given. Let $k30_abc Miz_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k32_abc Miz_1 : \iota \Rightarrow \iota$ be given. Let $k37_abc Miz_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_instalg1 X0) \wedge ((v1_abc Miz_1 X0) \wedge ((v3_abc Miz_1 \\ X0) \wedge (l1_msualg_1 X0)))) \Rightarrow (\forall X1.((\neg v6_abc Miz_1 X1 X0) \wedge (\\ m1_abc Miz_1 X1 X0 (k13_abc Miz_1 X0))) \Rightarrow (\exists X2.(m1_abc Miz_1 \\ X2 X0 (k13_abc Miz_1 X0)) \wedge ((X1 = k30_abc Miz_1 X0 (k32_abc Miz_1 X0 \\ X2) \wedge (k37_abc Miz_1 X0 X1 = X2)))))) \end{aligned} \tag{1}$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_instalg1 X0) \wedge ((v1_abc Miz_1 X0) \wedge ((v3_abc Miz_1 \\ X0) \wedge (l1_msualg_1 X0)))) \Rightarrow (\forall X1.((\neg v6_abc Miz_1 X1 X0) \wedge (\\ m1_abc Miz_1 X1 X0 (k13_abc Miz_1 X0))) \Rightarrow (k30_abc Miz_1 X0 (k32_abc Miz_1 \\ X0) (k37_abc Miz_1 X0 X1) = X1)) \end{aligned}$$