

t62_gfacirc1 (TMYKkWEaxaUwifw-
NaW8U9ARxR7zZTF7Ztgq)

October 27, 2020

Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_gfacirc1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_card_3 : \iota \Rightarrow \iota$ be given. Let $u3_msualg_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k22_gfacirc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k23_gfacirc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_margrel1 : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_twoscomp : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_facirc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k24_gfacirc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_margrel1 : \iota \Rightarrow \iota$ be given. Let $k2_binarith : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboolean : \iota \Rightarrow o$ be given. Let $k3_xboolean : \iota \Rightarrow \iota$ be given. Let $k10_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k3_xboolean (k10_xboolean X0 X1) = k10_xboolean (k3_xboolean X0) X1)) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(X2 \neq k4_tarski (k10_finseq_1 \\ & X0 X1) k4_gfacirc1) \Rightarrow (\forall X3.(m1_subset_1 X3 (k4_card_3 (u3_msualg_1 \\ & (k22_gfacirc1 X0 X1 X2) (k23_gfacirc1 X0 X1 X2)))) \Rightarrow (\forall X4. \\ & (m1_subset_1 X4 k6_margrel1) \Rightarrow (\forall X5.(m1_subset_1 X5 k6_margrel1) \Rightarrow \\ & (\forall X6.(m1_subset_1 X6 k6_margrel1) \Rightarrow (((X4 = k1_funct_1 X3 \\ & X0) \wedge ((X5 = k1_funct_1 X3 X1) \wedge (X6 = k1_funct_1 X3 X2))) \Rightarrow ((k1_twoscomp \\ & (k22_gfacirc1 X0 X1 X2) (k23_gfacirc1 X0 X1 X2) (k5_facirc_1 (k22_gfacirc1 \\ & X0 X1 X2) (k23_gfacirc1 X0 X1 X2) X3 np_2) (k24_gfacirc1 X0 X1 X2) = \\ & k2_binarith (k2_binarith X4 (k9_margrel1 X5)) (k9_margrel1 X6)) \wedge \\ & ((k1_funct_1 (k5_facirc_1 (k22_gfacirc1 X0 X1 X2) (k23_gfacirc1 \\ & X0 X1 X2) X3 np_2) (k4_tarski (k10_finseq_1 X0 X1) k4_gfacirc1) = \\ & k2_binarith X4 (k9_margrel1 X5)) \wedge ((k1_funct_1 (k5_facirc_1 (\\ & k22_gfacirc1 X0 X1 X2) (k23_gfacirc1 X0 X1 X2) X3 np_2) X0 = X4) \wedge (\\ & (k1_funct_1 (k5_facirc_1 (k22_gfacirc1 X0 X1 X2) (k23_gfacirc1 \\ & X0 X1 X2) X3 np_2) X1 = X5) \wedge (k1_funct_1 (k5_facirc_1 (k22_gfacirc1 \\ & X0 X1 X2) (k23_gfacirc1 X0 X1 X2) X3 np_2) X2 = X6)))))))))) \quad (2) \end{aligned}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k6_margrel1) \Rightarrow (k9_margrel1 X0 = k3_xboolean X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k6_margrel1) \wedge (m1_subset_1 X1 k6_margrel1)) \Rightarrow (k2_binarith X0 X1 = k10_xboolean X0 X1) \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k6_margrel1) \Rightarrow (m1_subset_1 (k9_margrel1 X0) k6_margrel1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k6_margrel1) \wedge (m1_subset_1 X1 k6_margrel1)) \Rightarrow (m1_subset_1 (k2_binarith X0 X1) k6_margrel1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.k4_tarski X0 X1 = k2_tarski (k2_tarski X0 X1) (k1_tarski X0) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (k10_xboolean X0 X1 = k10_xboolean X1 X0) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k6_margrel1) \Rightarrow (v1_xboolean X0) \quad (9)$$

Theorem 1

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(X2 \neq k4_tarski (k10_finseq_1 \\ & X0 X1) k4_gfacirc1) \Rightarrow (\forall X3.(m1_subset_1 X3 (k4_card_3 (u3_msualg_1 \\ & (k22_gfacirc1 X0 X1 X2) (k23_gfacirc1 X0 X1 X2)))) \Rightarrow (\forall X4. \\ & (m1_subset_1 X4 k6_margrel1) \Rightarrow (\forall X5.(m1_subset_1 X5 k6_margrel1) \Rightarrow \\ & (\forall X6.(m1_subset_1 X6 k6_margrel1) \Rightarrow (((X4 = k1_funct_1 X3 \\ & X0) \wedge ((X5 = k1_funct_1 X3 X1) \wedge (X6 = k1_funct_1 X3 X2))) \Rightarrow (k1_twoscomp \\ & (k22_gfacirc1 X0 X1 X2) (k23_gfacirc1 X0 X1 X2) (k5_facirc_1 (k22_gfacirc1 \\ & X0 X1 X2) (k23_gfacirc1 X0 X1 X2) X3 np_2) (k24_gfacirc1 X0 X1 X2) = \\ & k9_margrel1 (k2_binarith (k2_binarith X4 (k9_margrel1 X5)) X6)))))) \end{aligned}$$