

t62_interval1

(TMdsfo8imVaQzeSCH4ZJWky2oPqwg1TYvv6)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v3_roughs_1 : \iota \Rightarrow o$ be given. Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $m2_interval : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r2_interval : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k16_interval : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k15_interval : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k14_interval : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (((\neg v2_struct_0 X0) \wedge (v3_roughs_1 X0) \wedge (l1_orders_2 X0))) \wedge ((m2_interval X1 X0) \wedge (m2_interval X2 X0)) \Rightarrow ((r2_interval X0 X1 X2) \Leftrightarrow (X1 = X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (((\neg v2_struct_0 X0) \wedge (v3_roughs_1 X0) \wedge (l1_orders_2 X0))) \wedge ((m2_interval X1 X0) \wedge (m2_interval X2 X0)) \Rightarrow (m2_interval (k16_interval X0 X1 X2) X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (((\neg v2_struct_0 X0) \wedge (v3_roughs_1 X0) \wedge (l1_orders_2 X0))) \wedge (m2_interval X1 X0) \Rightarrow (m1_subset_1 (k15_interval X0 X1) (k1_zfmisc_1 (u1_struct_0 X0))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (((\neg v2_struct_0 X0) \wedge (v3_roughs_1 X0) \wedge (l1_orders_2 X0))) \wedge (m2_interval X1 X0) \Rightarrow (m1_subset_1 (k14_interval X0 X1) (k1_zfmisc_1 (u1_struct_0 X0))) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v3_roughs_1 X0) \wedge (l1_orders_2 \\ & X0))) \Rightarrow (\forall X1.(m2_interval X1 X0) \Rightarrow (\forall X2.(m2_interval \\ & X2 X0) \Rightarrow (k16_interval X0 X1 X2 = k4_tarski (k4_subset_1 (u1_struct_0 \\ & X0) (k14_interval X0 X1) (k14_interval X0 X2)) (k4_subset_1 (u1_struct_0 \\ & X0) (k15_interval X0 X1) (k15_interval X0 X2)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((m1_subset_1 X1 (k1_zfmisc_1 \\ & X0)) \wedge (m1_subset_1 X2 (k1_zfmisc_1 X0))) \Rightarrow (k4_subset_1 X0 X1 X2 = \\ & k4_subset_1 X0 X2 X1) \end{aligned} \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v3_roughs_1 X0) \wedge (l1_orders_2 \\ & X0))) \Rightarrow (\forall X1.(m2_interval X1 X0) \Rightarrow (\forall X2.(m2_interval \\ & X2 X0) \Rightarrow (r2_interval X0 (k16_interval X0 X1 X2) (k16_interval X0 \\ & X2 X1)))) \end{aligned}$$