

t62_xxreal_1

(TMd9E5tBvQZf45yhSs52atEY7NFcCrkPxWg)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((X0 \in k4_xxreal_1 X1 X2) \Leftrightarrow ((\neg r1_xxreal_0 X0 X1) \wedge \\ & (\neg r1_xxreal_0 X2 X0)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((X0 \in k1_xxreal_1 X1 X2) \Leftrightarrow ((r1_xxreal_0 X1 X0) \wedge \\ & (r1_xxreal_0 X0 X2)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (r1_xxreal_0 X0 X0) \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1) \Leftrightarrow (\forall X2.(X2 \in X0) \Rightarrow (X2 \in X1)) \tag{4}$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_tarski (k1_xxreal_1 X0 X1) (k4_xxreal_1 X2 X3))) \Rightarrow ((\\ & \neg r1_xxreal_0 X0 X2) \wedge (\neg r1_xxreal_0 X3 X1)))))) \end{aligned}$$