

t62_xxreal_2 (TMd-
NAXRx9w3qpx3WzbTEVw5eNjcxNGnUzpk)

October 27, 2020

Let $v2_membered : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xxreal_2 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v2_membered X1) \Rightarrow ((X0 \in X1) \Rightarrow (r1_xxreal_0 (k2_xxreal_2 X1) X0))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2.(v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow (r1_xxreal_0 X0 X2)))) \quad (2)$$

Assume the following.

$$\forall X0.(v2_membered X0) \Rightarrow (v1_xxreal_0 (k2_xxreal_2 X0)) \quad (3)$$

Theorem 1

$$\forall X0.(v2_membered X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((\exists X2.(v1_xxreal_0 X2) \wedge ((X2 \in X0) \wedge (r1_xxreal_0 X2 X1))) \Rightarrow (r1_xxreal_0 (k2_xxreal_2 X0) X1)))$$