

## t63\_cfunct\_1

(TMYW3Dcp98wTnPuutxonhAdgzsBpN5VG86J)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_numbers : \iota$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k8\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k1\_cfunct\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_cfunct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_valued\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k46\_valued\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k19\_valued\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_membered : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. ((v1\_funct\_1 X1) \wedge ( \\ & \quad m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k2\_numbers)))) \Rightarrow ( \\ & (v1\_partfun1 (k2\_cfunct\_1 X0 X1) X0) \Leftrightarrow ((k8\_relset\_1 X0 k2\_numbers \\ & \quad X1 (k1\_tarski k6\_numbers) = k1\_xboole\_0) \wedge (v1\_partfun1 X1 X0)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. ((v1\_funct\_1 X1) \wedge ( \\ & \quad m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k2\_numbers)))) \Rightarrow ( \\ & \quad \forall X2. ((v1\_funct\_1 X2) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & \quad X0 k2\_numbers)))) \Rightarrow (((v1\_partfun1 X1 X0) \wedge (v1\_partfun1 X2 X0)) \Rightarrow \\ & \quad (v1\_partfun1 (k2\_valued\_1 X0 k2\_numbers k2\_numbers X1 X2) X0)) \wedge \\ & \quad (((v1\_partfun1 (k2\_valued\_1 X0 k2\_numbers k2\_numbers X1 X2) X0) \Rightarrow \\ & \quad ((v1\_partfun1 X1 X0) \wedge (v1\_partfun1 X2 X0))) \wedge (((v1\_partfun1 X1 \\ & \quad X0) \wedge (v1\_partfun1 X2 X0)) \Rightarrow (v1\_partfun1 (k46\_valued\_1 X0 k2\_numbers \\ & \quad k2\_numbers X1 X2) X0)) \wedge (((v1\_partfun1 (k46\_valued\_1 X0 k2\_numbers \\ & \quad k2\_numbers X1 X2) X0) \Rightarrow ((v1\_partfun1 X1 X0) \wedge (v1\_partfun1 X2 X0))) \wedge \\ & \quad (((v1\_partfun1 X1 X0) \wedge (v1\_partfun1 X2 X0)) \Rightarrow (v1\_partfun1 (k19\_valued\_1 \\ & \quad X0 k2\_numbers k2\_numbers X1 X2) X0)) \wedge ((v1\_partfun1 (k19\_valued\_1 \\ & \quad X0 k2\_numbers k2\_numbers X1 X2) X0) \Rightarrow ((v1\_partfun1 X1 X0) \wedge (v1\_partfun1 \\ & \quad X2 X0)))))))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge ( \\ m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k2\_numbers)))) \Rightarrow ( \\ \forall X2.((v1\_funct\_1 X2) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ X0 k2\_numbers)))) \Rightarrow (r2\_relset\_1 X0 k2\_numbers (k1\_cfunct\_1 X0 \\ X1 X2) (k19\_valued\_1 X0 k2\_numbers k2\_numbers X1 (k2\_cfunct\_1 X0 \\ X2)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.\forall X3.((m1\_subset\_1 X2 \\ (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 X1)))) \Rightarrow ((r2\_relset\_1 X0 X1 X2 X3) \Leftrightarrow (X2 = X3)) \end{aligned} \quad (4)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (5)$$

Assume the following.

$$v1\_membered k2\_numbers \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((v1\_funct\_1 X1) \wedge ( \\ m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k2\_numbers)))) \Rightarrow ( \\ ((v1\_funct\_1 (k2\_cfunct\_1 X0 X1)) \wedge (m1\_subset\_1 (k2\_cfunct\_1 \\ X0 X1) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k2\_numbers)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0) \wedge (((v1\_funct\_1 \\ X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k2\_numbers)))) \wedge \\ ((v1\_funct\_1 X2) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 \\ k2\_numbers)))))) \Rightarrow ((v1\_funct\_1 (k1\_cfunct\_1 X0 X1 X2)) \wedge (m1\_subset\_1 \\ (k1\_cfunct\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k2\_numbers)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.\forall X3.\forall X4.((v1\_membered \\ X1) \wedge ((v1\_membered X2) \wedge (((v1\_funct\_1 X3) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 X1)))) \wedge ((v1\_funct\_1 X4) \wedge (m1\_subset\_1 X4 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 X2)))))) \Rightarrow ((v1\_funct\_1 (k19\_valued\_1 X0 X1 X2 \\ X3 X4)) \wedge (m1\_subset\_1 (k19\_valued\_1 X0 X1 X2 X3 X4) (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 k2\_numbers)))))) \end{aligned} \quad (9)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge ( \\ m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k2\_numbers)))) \Rightarrow ( \\ \forall X2.((v1\_funct\_1 X2) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ X0 k2\_numbers)))) \Rightarrow (((v1\_partfun1 X1 X0) \wedge ((k8\_relset\_1 X0 k2\_numbers \\ X2 (k1\_tarski k6\_numbers) = k1\_xboole\_0) \wedge (v1\_partfun1 X2 X0))) \Leftrightarrow \\ (v1\_partfun1 (k1\_cfunct\_1 X0 X1 X2) X0)))) \end{aligned}$$