

t63_fvaluat1 (TM- dugxDESkGb7BGLpV4hgMphwvT62y1kxB7)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v6_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v3_group_1 : \iota \Rightarrow o$ be given. Let $v5_vectsp_1 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v1_realset2 : \iota \Rightarrow o$ be given. Let $l6_algstr_0 : \iota \Rightarrow o$ be given. Let $m1_fvaluat1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_fvaluat1 : \iota \Rightarrow o$ be given. Let $k5_struct_0 : \iota \Rightarrow \iota$ be given. Let $k8_fvaluat1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_fvaluat1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((\neg v6_struct_0 X0) \wedge ((v13_algstr_0 \\ & X0) \wedge ((v3_group_1 X0) \wedge ((v5_vectsp_1 X0) \wedge ((v2_rlvect_1 X0) \wedge (\\ & (v3_rlvect_1 X0) \wedge (v4_rlvect_1 X0) \wedge (v1_realset2 X0) \wedge (l6_algstr_0 \\ & X0)))))))))) \Rightarrow (\forall X1. (m1_fvaluat1 X1 X0) \Rightarrow ((v3_fvaluat1 \\ & X0) \Rightarrow (k5_struct_0 X0 \in k6_fvaluat1 X0 X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge (l6_algstr_0 X0)) \Rightarrow (\forall X1. \\ & (m1_fvaluat1 X1 X0) \Rightarrow (\forall X2. (m1_subset_1 X2 (u1_struct_0 \\ & X0)) \Rightarrow ((X2 \in k6_fvaluat1 X0 X1) \Leftrightarrow (r1_xxreal_0 k6_numbers (k3_funct_2 \\ & (u1_struct_0 X0) k7_numbers X1 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((\neg v6_struct_0 X0) \wedge ((v13_algstr_0 \\ & X0) \wedge ((v3_group_1 X0) \wedge ((v5_vectsp_1 X0) \wedge ((v2_rlvect_1 X0) \wedge (\\ & (v3_rlvect_1 X0) \wedge (v4_rlvect_1 X0) \wedge (v1_realset2 X0) \wedge (l6_algstr_0 \\ & X0)))))))))) \Rightarrow (\forall X1. (m1_fvaluat1 X1 X0) \Rightarrow ((v3_fvaluat1 \\ & X0) \Rightarrow (k3_funct_2 (u1_struct_0 X0) k7_numbers X1 (k5_struct_0 X0) = \\ & k6_numbers))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge ((\neg v6_struct_0 X0) \wedge ((v13_algstr_0 \\
& X0) \wedge ((v3_group_1 X0) \wedge ((v5_vectsp_1 X0) \wedge ((v2_rlvect_1 X0) \wedge \\
& (v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v1_realset2 X0) \wedge (l6_algstr_0 \\
& X0)))))))))) \Rightarrow (\forall X1.(m1_fvaluat1 X1 X0) \Rightarrow ((v3_fvaluat1 \\
& X0) \Rightarrow (k8_fvaluat1 X0 X1 = ReplSep (toset (\lambda X2 : \iota.m1_subset_1 \\
& X2 (u1_struct_0 X0))) (\lambda X2 : \iota.\neg r1_xxreal_0 (k3_funct_2 (\\
& u1_struct_0 X0) k7_numbers X1 X2) k6_numbers) (\lambda X2 : \iota.X2))))))
\end{aligned} \tag{4}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge ((\neg v6_struct_0 X0) \wedge ((v13_algstr_0 \\
& X0) \wedge ((v3_group_1 X0) \wedge ((v5_vectsp_1 X0) \wedge ((v2_rlvect_1 X0) \wedge \\
& (v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v1_realset2 X0) \wedge (l6_algstr_0 \\
& X0)))))))))) \Rightarrow (\forall X1.(m1_fvaluat1 X1 X0) \Rightarrow (\neg (v3_fvaluat1 \\
& X0) \wedge (k5_struct_0 X0 \in k8_fvaluat1 X0 X1)))
\end{aligned}$$