

t63_sprect_1 (TMT- EDgU3u9bYLbqnpjEZKp7QCtwkjyXroJW)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v2_compts_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k12_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k3_topreal1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_sprect_1 : \iota \Rightarrow \iota$ be given. Let $k8_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k7_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0. (& \neg v1_xboole_0 X0) \wedge ((v2_compts_1 X0 (k15_euclid np_2)) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))))) \Rightarrow \\ & (k8_pscomp_1 (k3_topreal1 np_2 (k1_sprect_1 X0)) = k8_pscomp_1 \\ & X0) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. (& \neg v1_xboole_0 X0) \wedge ((v2_compts_1 X0 (k15_euclid np_2)) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))))) \Rightarrow \\ & (k7_pscomp_1 (k3_topreal1 np_2 (k1_sprect_1 X0)) = k7_pscomp_1 \\ & X0) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (4)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(m1_finseq_1\ X1\ (u1_struct_0\ (k15_euclid\ X0))))\Rightarrow(m1_subset_1\ (k3_topreal1\ X0\ X1)\ (k1_zfmisc_1\ (u1_struct_0\ (k15_euclid\ X0)))) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ (k1_zfmisc_1\ (u1_struct_0\ (k15_euclid\ np_2))))\Rightarrow(m2_finseq_1\ (k1_sprect_1\ X0)\ (u1_struct_0\ (k15_euclid\ np_2))) \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ (k1_zfmisc_1\ (u1_struct_0\ (k15_euclid\ np_2))))\Rightarrow(k12_pscomp_1\ X0 = k19_euclid\ (k8_pscomp_1\ X0)\ (k7_pscomp_1\ X0)) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k4_ordinal1)\Rightarrow(v7_ordinal1\ X0) \quad (9)$$

Theorem 1

$$\forall X0.((\neg v1_xboole_0\ X0)\wedge((v2_compts_1\ X0\ (k15_euclid\ np_2))\wedge(m1_subset_1\ X0\ (k1_zfmisc_1\ (u1_struct_0\ (k15_euclid\ np_2))))))\Rightarrow(k12_pscomp_1\ (k3_topreal1\ np_2\ (k1_sprect_1\ X0)) = k12_pscomp_1\ X0)$$