

# t63\_topreal6 (TMRqHiPTgAwqBRSUXP- sHDxFMdDDw7UdSUHB)

October 27, 2020

Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k15\_euclid : \iota \Rightarrow \iota$  be given. Let  $v9\_rltopsp1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_pre\_topc : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v6\_membered : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v2\_pre\_topc : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v6\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v7\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v8\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v6\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $v7\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $l1\_rltopsp1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2. (m2\_subset\_1 \\ & X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \quad (1)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (2)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1) \wedge (v3\_ordinal1 k4\_ordinal1) \quad (3)$$

Assume the following.

$$v6\_membered k4\_ordinal1 \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. (v7\_ordinal1 X0) \Rightarrow ((v2\_pre\_topc (k15\_euclid X0)) \wedge \\ & ((v13\_algstr\_0 (k15\_euclid X0)) \wedge ((v2\_rlvect\_1 (k15\_euclid X0)) \wedge \\ & ((v3\_rlvect\_1 (k15\_euclid X0)) \wedge ((v4\_rlvect\_1 (k15\_euclid X0)) \wedge \\ & ((v5\_rlvect\_1 (k15\_euclid X0)) \wedge ((v6\_rlvect\_1 (k15\_euclid X0)) \wedge \\ & ((v7\_rlvect\_1 (k15\_euclid X0)) \wedge ((v8\_rlvect\_1 (k15\_euclid X0)) \wedge \\ & (v5\_rltopsp1 (k15\_euclid X0)))))))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow ((\neg v2\_struct\_0 (k15\_euclid X0)) \wedge (v5\_rltopsp1 (k15\_euclid X0))) \quad (6)$$

Assume the following.

$$\neg v1\_xboole\_0 k1\_numbers \quad (7)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow ((v5\_rltopsp1 (k15\_euclid X0)) \wedge ((v6\_rltopsp1 (k15\_euclid X0)) \wedge (v7\_rltopsp1 (k15\_euclid X0)))) \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(((\neg v2\_struct\_0 X0) \wedge (v2\_pre\_topc X0) \wedge \\ & ((v13\_algstr\_0 X0) \wedge (v2\_rlvect\_1 X0) \wedge (v3\_rlvect\_1 X0) \wedge (v4\_rlvect\_1 \\ & X0) \wedge (v5\_rlvect\_1 X0) \wedge (v6\_rlvect\_1 X0) \wedge (v7\_rlvect\_1 X0) \wedge \\ & ((v8\_rlvect\_1 X0) \wedge (v6\_rltopsp1 X0) \wedge (v7\_rltopsp1 X0) \wedge (l1\_rltopsp1 \\ & X0)))))) \wedge ((v9\_rltopsp1 X1 X0) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 \\ & (u1\_struct\_0 X0)))) \Rightarrow (v9\_rltopsp1 (k2\_pre\_topc X0 X1) X0) \end{aligned} \quad (9)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (10)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow ((v5\_rltopsp1 (k15\_euclid X0)) \wedge (l1\_rltopsp1 (k15\_euclid X0))) \quad (11)$$

Assume the following.

$$\forall X0.(v6\_membered X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 X0) \Rightarrow (v7\_ordinal1 X1)) \quad (12)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(m2\_subset\_1 X0 k1\_numbers k5\_numbers) \Rightarrow (\forall X1. \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 (k15\_euclid X0)))) \Rightarrow \\ & ((v9\_rltopsp1 X1 (k15\_euclid X0)) \Rightarrow (v9\_rltopsp1 (k2\_pre\_topc \\ & (k15\_euclid X0) X1) (k15\_euclid X0)))) \end{aligned}$$