

t63_waybel34
(TMHqXjy4beebg7nE1o1G14VqNsknngKHtiT)

October 27, 2020

Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_yellow_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $u1_orders_2 : \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow ((r1_tarski X0 (k9_xtuple_0 X1)) \Rightarrow (k9_xtuple_0 (k5_relat_1 X1 X0) = X0)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_tarski (k10_xtuple_0 X1) X0) \Rightarrow ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 (k9_xtuple_0 X1) X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 (k9_xtuple_0 X1) X0)))))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k10_xtuple_0 (k5_relat_1 X1 X0) = k7_relat_1 X1 X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (k7_relset_1 X0 X1 X2 X3 = k7_relat_1 X2 X3) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow(k2_partfun1 X0 X1 X2 X3 = k5_relat_1 X2 X3) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (6)$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0)\wedge(l1_struct_0 X0))\Rightarrow(\neg v1_xboole_0 (u1_struct_0 X0)) \quad (7)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (8)$$

Assume the following.

$$\forall X0.(l1_orders_2 X0)\Rightarrow(\forall X1.(m1_yellow_0 X1 X0)\Rightarrow (l1_orders_2 X1)) \quad (9)$$

Assume the following.

$$\forall X0.(l1_orders_2 X0)\Rightarrow(l1_struct_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((v1_funct_1 (k2_partfun1 X0 X1 X2 X3))\wedge(m1_subset_1 (k2_partfun1 X0 X1 X2 X3) (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(((X1\neq k1_xboole_0)\Rightarrow((v1_funct_2 X2 X0 X1)\Leftrightarrow(X0 = k1_relset_1 X0 X2)))\wedge((X1 = k1_xboole_0)\Rightarrow((v1_funct_2 X2 X0 X1)\Leftrightarrow(X2 = k1_xboole_0)))) \quad (12)$$

Assume the following.

$$\forall X0.(l1_orders_2 X0)\Rightarrow(\forall X1.(l1_orders_2 X1)\Rightarrow((m1_yellow_0 X1 X0)\Leftrightarrow((r1_tarski (u1_struct_0 X1) (u1_struct_0 X0))\wedge(r1_tarski (u1_orders_2 X1) (u1_orders_2 X0)))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (15)$$

Theorem 1

$$\begin{aligned} & \forall X0.(l1_orders_2 X0)\Rightarrow(\forall X1.((\neg v2_struct_0 X1)\wedge \\ & (l1_orders_2 X1))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge((v1_funct_2 \\ & X2 (u1_struct_0 X0) (u1_struct_0 X1))\wedge(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 X1))))))\Rightarrow(\forall X3. \\ & (m1_yellow_0 X3 X0)\Rightarrow(\forall X4.(m1_yellow_0 X4 X1)\Rightarrow((r1_tarski \\ & (k7_reset_1 (u1_struct_0 X0) (u1_struct_0 X1) X2 (u1_struct_0 \\ & X3)) (u1_struct_0 X4))\Rightarrow((v1_funct_1 (k2_partfun1 (u1_struct_0 \\ & X0) (u1_struct_0 X1) X2 (u1_struct_0 X3)))\wedge((v1_funct_2 (k2_partfun1 \\ & (u1_struct_0 X0) (u1_struct_0 X1) X2 (u1_struct_0 X3)) (u1_struct_0 \\ & X3) (u1_struct_0 X4))\wedge(m1_subset_1 (k2_partfun1 (u1_struct_0 \\ & X0) (u1_struct_0 X1) X2 (u1_struct_0 X3)) (k1_zfmisc_1 (k2_zfmisc_1 \\ & (u1_struct_0 X3) (u1_struct_0 X4))))))))))))) \end{aligned}$$