

t64\_funct\_4  
(TMT5BRn9CtxoiPqFbVSaJC71HL9m3Qm8h9L)

October 27, 2020

Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k4\_funct\_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((X0 \neq X1) \Rightarrow (k1\_funct\_1 \\ & (k4\_funct\_4 X0 X1 X2 X3) X0 = X2)) \wedge (k1\_funct\_1 (k4\_funct\_4 X0 X1 X2 \\ & X3) X1 = X3) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (k9\_xtuple\_0 (k4\_funct\_4 \\ & X0 X1 X2 X3) = k2\_tarski X0 X1) \wedge (r1\_tarski (k10\_xtuple\_0 (k4\_funct\_4 \\ & X0 X1 X2 X3)) (k2\_tarski X2 X3)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (v1\_relat\_1 (k4\_funct\_4 \\ & X0 X1 X2 X3)) \wedge (v1\_funct\_1 (k4\_funct\_4 X0 X1 X2 X3)) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (r1\_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow \\ & (X2 \in X1)) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. (X1 = \\ & k10\_xtuple\_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X3 \in k9\_xtuple\_0 \\ & X0) \wedge (X2 = k1\_funct\_1 X0 X3)))) \end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (X2 = k2\_tarski X0 X1) \Leftrightarrow (\forall X3. \\ & (X3 \in X2) \Leftrightarrow ((X3 = X0) \vee (X3 = X1))) \end{aligned} \tag{6}$$

Assume the following.

$$\forall X0.\forall X1.(X0 = X1) \Leftrightarrow ((r1\_tarSKI X0 X1) \wedge (r1\_tarSKI X1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.k2\_tarSKI X0 X1 = k2\_tarSKI X1 X0 \quad (8)$$

**Theorem 1**

$$\forall X0.\forall X1.\forall X2.\forall X3.(X0 \neq X1) \Rightarrow (k10\_xtuple\_0 \\ (k4\_funct\_4 X0 X1 X2 X3) = k2\_tarSKI X2 X3)$$