

t64_fvaluat1

(TMbR3mAQJKrDEFv5yURdv5yxSgjETxy9YpA)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v6_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v3_group_1 : \iota \Rightarrow o$ be given. Let $v5_vectsp_1 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v1_realset2 : \iota \Rightarrow o$ be given. Let $l6_algstr_0 : \iota \Rightarrow o$ be given. Let $m1_fvaluat1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v3_fvaluat1 : \iota \Rightarrow o$ be given. Let $k7_fvaluat1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_fvaluat1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $k13_ideal_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $v2_fvaluat1 : \iota \Rightarrow o$ be given. Let $l2_algstr_0 : \iota \Rightarrow o$ be given. Let $l5_algstr_0 : \iota \Rightarrow o$ be given. Let $l2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_algstr_0 : \iota \Rightarrow o$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_supinf_2 : \iota \Rightarrow \iota$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. r1_tarski (k3_xboole_0 X0 X1) X0 \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (k8_relset_1 X0 X1 X2 X3 = k8_relat_1 X2 X3) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))))) \Rightarrow (k13_ideal_1 X0 X1 X2 = k3_xboole_0 X1 X2) \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.(l6_algstr_0 X0) \Rightarrow (\forall X1.(m1_fvaluat1 X1 X0) \Rightarrow \\ & ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 (u1_struct_0 X0) k7_numbers) \wedge \\ & ((v2_fvaluat1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & (u1_struct_0 X0) k7_numbers))))))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.(l6_algstr_0 X0) \Rightarrow ((l2_algstr_0 X0) \wedge (l5_algstr_0 X0)) \quad (5)$$

Assume the following.

$$\forall X0.(l2_struct_0 X0) \Rightarrow (l1_struct_0 X0) \quad (6)$$

Assume the following.

$$\forall X0.(l2_algstr_0 X0) \Rightarrow ((l2_struct_0 X0) \wedge (l1_algstr_0 X0)) \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (\\ & k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (m1_subset_1 (k8_relset_1 \\ & X0 X1 X2 X3) (k1_zfmisc_1 X0)) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((\neg v6_struct_0 X0) \wedge ((v13_algstr_0 \\ & X0) \wedge ((v3_group_1 X0) \wedge ((v5_vectsp_1 X0) \wedge ((v2_rlvect_1 X0) \wedge (\\ & (v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v1_realset2 X0) \wedge (l6_algstr_0 \\ & X0)))))))))) \Rightarrow (\forall X1.(m1_fvaluat1 X1 X0) \Rightarrow (\forall X2.((\\ & \neg v1_xboole_0 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow \\ & (((v3_fvaluat1 X0) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 \\ & (k7_fvaluat1 X0 X1)))))) \Rightarrow (k9_fvaluat1 X0 X1 X2 = k13_ideal_1 X0 (\\ & k8_relset_1 (u1_struct_0 X0) k7_numbers X1 (k6_domain_1 k7_numbers \\ & (k7_supinf_2 (k7_relset_1 (u1_struct_0 X0) k7_numbers X1 X2)))) \\ & X2)))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (10)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((\neg v6_struct_0 X0) \wedge ((v13_algstr_0 \\ & X0) \wedge ((v3_group_1 X0) \wedge ((v5_vectsp_1 X0) \wedge ((v2_rlvect_1 X0) \wedge (\\ & (v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v1_realset2 X0) \wedge (l6_algstr_0 \\ & X0)))))))))) \Rightarrow (\forall X1.(m1_fvaluat1 X1 X0) \Rightarrow (\forall X2.((\\ & \neg v1_xboole_0 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow \\ & (((v3_fvaluat1 X0) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 \\ & (k7_fvaluat1 X0 X1)))))) \Rightarrow (r1_tarski (k9_fvaluat1 X0 X1 X2) X2)))) \end{aligned}$$