

t64_lukasi_1

(TMZPcpvPAdTskdArhFEXa9ajrcJ8keewLVY)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r3_cqc_the1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X3.(m2_subset_1 X3 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X4.(m1_subset_1 X4 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow ((r3_cqc_the1 X0 X4 (k8_cqc_lang X0 X1 (k8_cqc_lang \\ X0 X2 X3))) \Rightarrow (r3_cqc_the1 X0 X4 (k8_cqc_lang X0 X2 (k8_cqc_lang X0 \\ X1 X3)))))))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow (\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X3.(m2_subset_1 X3 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (((r3_cqc_the1 X0 X1 X2) \wedge (r3_cqc_the1 X0 \\ X1 (k8_cqc_lang X0 X2 X3))) \Rightarrow (r3_cqc_the1 X0 X1 X3)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (\neg v1_xboole_0 (k3_cqc_lang X0)) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((m1_qc_lang1 X0) \wedge ((m1_subset_1 \\ X1 (k3_cqc_lang X0)) \wedge (m1_subset_1 X2 (k3_cqc_lang X0)))) \Rightarrow (m2_subset_1 \\ (k8_cqc_lang X0 X1 X2) (k9_qc_lang1 X0) (k3_cqc_lang X0)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(m1_qc_lang1\ X0)\Rightarrow(m1_subset_1\ (k3_cqc_lang\ X0)\ (k1_zfmisc_1\ (k9_qc_lang1\ X0))) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xboole_0\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ X0))\Rightarrow(v1_xboole_0\ X1)) \quad (7)$$

Theorem 1

$$\begin{aligned} &\forall X0.(m1_qc_lang1\ X0)\Rightarrow(\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1\ X0)\ (k3_cqc_lang\ X0))\Rightarrow(\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1\ X0)\ (k3_cqc_lang\ X0))\Rightarrow(\forall X3.(m2_subset_1\ X3\ (k9_qc_lang1\ X0)\ (k3_cqc_lang\ X0))\Rightarrow(\forall X4.(m1_subset_1\ X4\ (k1_zfmisc_1\ (k3_cqc_lang\ X0))\Rightarrow(((r3_cqc_the1\ X0\ X4\ (k8_cqc_lang\ X0\ X1\ (k8_cqc_lang\ X0\ X2\ X3)))\wedge(r3_cqc_the1\ X0\ X4\ X2))\Rightarrow(r3_cqc_the1\ X0\ X4\ (k8_cqc_lang\ X0\ X1\ X3)))))))))) \end{aligned}$$