

t65_arytm_3 (TMbETJKUy-
LufF3qj5wZdmmQFnSNGhCmhRFT)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_arytm_3 : \iota$ be given. Let $r3_arytm_3 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k11_arytm_3 : \iota$ be given. Let $k9_arytm_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(m1_subset_1 X0 k5_arytm_3) \Rightarrow (\forall X1.(m1_subset_1 X1 k5_arytm_3) \Rightarrow ((k9_arytm_3 X0 X1 = k11_arytm_3) \Rightarrow (X0 = k11_arytm_3))) \quad (1)$$

Assume the following.

$$m1_subset_1 k11_arytm_3 k5_arytm_3 \quad (2)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k5_arytm_3) \Rightarrow (\forall X1.(m1_subset_1 X1 k5_arytm_3) \Rightarrow ((r3_arytm_3 X0 X1) \Leftrightarrow (\exists X2.(m1_subset_1 X2 k5_arytm_3) \wedge (X1 = k9_arytm_3 X0 X2)))) \quad (3)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k5_arytm_3) \Rightarrow ((r3_arytm_3 X0 k11_arytm_3) \Rightarrow (X0 = k11_arytm_3))$$