

# t65\_bvfunc\_1

(TMU3fViLWrzTPQkCkzBrh8j4Hr6rnkgkURU)

October 27, 2020

Let  $v1\_xboolean : \iota \Rightarrow o$  be given. Let  $k8\_margrel1 : \iota$  be given. Let  $k6\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xboolean : \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (( \\ (k6\_xboolean X0 X1 = k2\_xboolean) \wedge (k6\_xboolean X1 X0 = k2\_xboolean)) \Rightarrow \\ (X0 = X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (k6\_xboolean \\ X0 (k6\_xboolean X1 X0) = k2\_xboolean)) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xboolean X0) \Rightarrow (k6\_xboolean X0 X0 = k2\_xboolean) \quad (3)$$

Assume the following.

$$k8\_margrel1 = k2\_xboolean \quad (4)$$

## Theorem 1

$$\begin{aligned} \forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (( \\ (X0 = k8\_margrel1) \wedge (k6\_xboolean X0 X1 = k8\_margrel1)) \Rightarrow (X1 = k8\_margrel1))) \end{aligned}$$