

t65_complfld

(TMcy1MCzbvGtHd7iSEcuv6ZYLDC2pTj1Zmr)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_complfld : \iota$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k17_complex1 : \iota \Rightarrow \iota$ be given. Let $k5_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k9_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k16_complex1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (r1_xreal_0 \\ (k9_real_1 (k17_complex1 X0) (k17_complex1 X1)) (k17_complex1 \\ (k6_xcmplx_0 X0 X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (u1_struct_0 k1_complfld)) \Rightarrow (\forall X1. \\ (m1_subset_1 X1 (u1_struct_0 k1_complfld)) \Rightarrow (\forall X2.(v1_xcmplx_0 \\ X2) \Rightarrow (\forall X3.(v1_xcmplx_0 X3) \Rightarrow (((X0 = X2) \wedge (X1 = X3)) \Rightarrow (k5_algstr_0 \\ k1_complfld X0 X1 = k4_binop_2 X2 X3)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (k9_real_1 X0 X1 = k6_xcmplx_0 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k4_binop_2 X0 X1 = k6_xcmplx_0 X0 X1) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k17_complex1 X0 = k16_complex1 X0) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (k10_binop_2 X0 X1 = k6_xcmplx_0 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (m1_subset_1 (k17_complex1 X0) k1_numbers) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (v1_xreal_0 (k16_complex1 X0)) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 k1_complfld)) \Rightarrow (v1_xcmplx_0 X0) \quad (9)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 k1_complfld)) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 k1_complfld)) \Rightarrow (r1_xreal_0 (k10_binop_2 (k17_complex1 X0) (k17_complex1 X1)) (k17_complex1 (k5_algstr_0 k1_complfld X0 X1))))$$