

t65\_gfacirc1  
(TMRaGx9P8DWjobY5xqffj8a6puBfaU7T938)

October 27, 2020

Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k10\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_gfacirc1 : \iota$  be given. Let  $k3\_twoscomp : \iota$  be given. Let  $k2\_twoscomp : \iota$  be given. Let  $k3\_gfacirc1 : \iota$  be given. Let  $k2\_msafree2 : \iota \Rightarrow \iota$  be given. Let  $k25\_gfacirc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $k6\_margrel1 : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k8\_facirc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v1\_circcomb : \iota \Rightarrow o$  be given. Let  $v2\_circcomb : \iota \Rightarrow o$  be given. Let  $l1\_msualg\_1 : \iota \Rightarrow o$  be given. Let  $r1\_circcomb : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k19\_gfacirc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_circcomb : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k22\_gfacirc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v11\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v1\_msualg\_1 : \iota \Rightarrow o$  be given. Let  $v3\_circcomb : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1\_funct\_1 X3) \wedge \\ & ((v1\_funct\_2 X3 (k4\_finseq\_2 np\_2 k6\_margrel1) k6\_margrel1) \wedge \\ & (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (k4\_finseq\_2 np\_2 \\ & k6\_margrel1) k6\_margrel1)))))) \Rightarrow ((X0 \neq k4\_tarski (k10\_finseq\_1 \\ & X1 X2) X3) \Rightarrow (k2\_msafree2 (k8\_facirc\_1 X1 X2 X0 X3) = k1\_enumset1 X1 \\ & X2 X0)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v1\_circcomb X0) \wedge ((v2\_circcomb \\ & X0) \wedge (l1\_msualg\_1 X0)))) \Rightarrow (\forall X1. ((\neg v2\_struct\_0 X1) \wedge ((v1\_circcomb \\ & X1) \wedge ((v2\_circcomb X1) \wedge (l1\_msualg\_1 X1)))) \Rightarrow (r1\_circcomb X0 X1)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \neg (X0 \neq k4\_tarski (k10\_finseq\_1 \\ & X1 X2) k3\_twoscomp) \wedge ((X1 \neq k4\_tarski (k10\_finseq\_1 X2 X0) k2\_twoscomp) \wedge \\ & ((X2 \neq k4\_tarski (k10\_finseq\_1 X0 X1) k3\_gfacirc1) \wedge (k2\_msafree2 \\ & (k19\_gfacirc1 X0 X1 X2) \neq k1\_enumset1 X0 X1 X2))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge (l1\_msualg\_1 X0)) \Rightarrow (\forall X1. \\ & ((\neg v2\_struct\_0 X1) \wedge (l1\_msualg\_1 X1)) \Rightarrow (((r1\_circcomb X0 X1) \wedge \\ & (k2\_msafree2 X0 = k2\_msafree2 X1)) \Rightarrow (k2\_msafree2 (k2\_circcomb \\ & X0 X1) = k2\_msafree2 X0))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & (v1\_funct\_1 k4\_gfacirc1) \wedge ((v1\_funct\_2 k4\_gfacirc1 (k4\_finseq\_2 \\ & np\_2 k6\_margrel1) k6\_margrel1) \wedge (m1\_subset\_1 k4\_gfacirc1 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 (k4\_finseq\_2 np\_2 k6\_margrel1) k6\_margrel1)))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (\neg v2\_struct\_0 (k22\_gfacirc1 \\ & X0 X1 X2)) \wedge ((\neg v11\_struct\_0 (k22\_gfacirc1 X0 X1 X2)) \wedge ((v1\_msualg\_1 \\ & (k22\_gfacirc1 X0 X1 X2)) \wedge ((v1\_circcomb (k22\_gfacirc1 X0 X1 X2)) \wedge \\ & ((v2\_circcomb (k22\_gfacirc1 X0 X1 X2)) \wedge ((v3\_circcomb (k22\_gfacirc1 \\ & X0 X1 X2)) \wedge (l1\_msualg\_1 (k22\_gfacirc1 X0 X1 X2))))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (\neg v2\_struct\_0 (k19\_gfacirc1 \\ & X0 X1 X2)) \wedge ((\neg v11\_struct\_0 (k19\_gfacirc1 X0 X1 X2)) \wedge ((v1\_msualg\_1 \\ & (k19\_gfacirc1 X0 X1 X2)) \wedge ((v1\_circcomb (k19\_gfacirc1 X0 X1 X2)) \wedge \\ & ((v2\_circcomb (k19\_gfacirc1 X0 X1 X2)) \wedge ((v3\_circcomb (k19\_gfacirc1 \\ & X0 X1 X2)) \wedge (l1\_msualg\_1 (k19\_gfacirc1 X0 X1 X2))))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. k25\_gfacirc1 X0 X1 X2 = k2\_circcomb \\ & (k22\_gfacirc1 X0 X1 X2) (k19\_gfacirc1 X0 X1 X2) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. k22\_gfacirc1 X0 X1 X2 = k8\_facirc1 \\ & X0 X1 X2 k4\_gfacirc1 \end{aligned} \quad (9)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \neg (X2 \neq k4\_tarski (k10\_finseq\_1 \\ & X0 X1) k4\_gfacirc1) \wedge ((X0 \neq k4\_tarski (k10\_finseq\_1 X1 X2) k3\_twoscomp) \wedge \\ & ((X1 \neq k4\_tarski (k10\_finseq\_1 X2 X0) k2\_twoscomp) \wedge ((X2 \neq k4\_tarski \\ & (k10\_finseq\_1 X0 X1) k3\_gfacirc1) \wedge (k2\_msafree2 (k25\_gfacirc1 \\ & X0 X1 X2) \neq k1\_enumset1 X0 X1 X2)))) \end{aligned}$$