

t65_pepin
(TMUa1Qq8BWZcLvZ6P3RRdzPjMy7QhkPhmPV)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_power : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (k3_power X0 X1 = k1_newton X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 np_1) \wedge ((\neg r1_xxreal_0 X1 k6_numbers) \wedge (r1_xxreal_0 (k3_power X0 X1) np_1)))) \quad (2)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (3)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xreal_0 X0) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X1 k6_numbers) \wedge ((\neg r1_xxreal_0 X0 np_1) \wedge (r1_xxreal_0 (k1_newton X0 X1) np_1))))$$