

t65\_xcplx\_1  
(TMJoftHsFi4Mwmey8UPfleSxVdjQcwBeruf)

October 27, 2020

Let  $v1\_xcplx\_0 : \iota \Rightarrow o$  be given. Let  $k2\_xcplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_xcplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xcplx\_0 X0) \Rightarrow (k7\_xcplx\_0 (k2\_xcplx\_0 X0 X0) \quad np\_2 = X0) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xcplx\_0 X0) \Rightarrow (\forall X1.(v1\_xcplx\_0 X1) \Rightarrow (\forall X2.(v1\_xcplx\_0 X2) \Rightarrow (k2\_xcplx\_0 (k7\_xcplx\_0 X0 X1) (k7\_xcplx\_0 X2 X1) = k7\_xcplx\_0 (k2\_xcplx\_0 X0 X2) X1))) \quad (2)$$

Assume the following.

$$((v2\_xxreal\_0 np\_2) \wedge (m2\_subset\_1 np\_2 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_2 k5\_numbers) \wedge (m1\_subset\_1 np\_2 k1\_numbers)) \quad (3)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xcplx\_0 X0) \quad (4)$$

**Theorem 1**

$$\forall X0.(v1\_xcplx\_0 X0) \Rightarrow (k2\_xcplx\_0 (k7\_xcplx\_0 X0 np\_2) (k7\_xcplx\_0 X0 np\_2) = X0)$$