

t66_finseq_6

(TMUTu7UdRF2ipo1dzjGB8c3smR1jMUMek8U)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_finseq_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow \\ & (\forall X2. (m2_finseq_1 X2 X0) \Rightarrow ((X1 \in k10_xtuple_0 X2) \Rightarrow (k1_finseq_5 \\ & X0 X2 X1 = k7_finseq_1 (k5_finseq_4 X2 X1) (k12_finseq_1 X0 X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1)))) \Rightarrow (k10_xtuple_0 (k7_finseq_1 X0 X1) = k2_xboole_0 (k10_xtuple_0 \\ & X0) (k10_xtuple_0 X1)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1)))) \Rightarrow (\forall X2. (X2 \in k10_xtuple_0 X0) \Rightarrow (k5_finseq_4 (k7_finseq_1 \\ & X0 X1) X2 = k5_finseq_4 X0 X2)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1_finseq_1 X1 X0) \wedge (m1_finseq_1 X2 X0)) \Rightarrow (k8_finseq_1 X0 X1 X2 = k7_finseq_1 X1 X2) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_finseq_1 X1 X0)\wedge(m1_finseq_1 X2 X0))\Rightarrow(m2_finseq_1 (k8_finseq_1 X0 X1 X2) X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k2_xboole_0 X0 X1)\Leftrightarrow(\forall X3.(X3 \in X2)\Leftrightarrow((X3 \in X0)\vee(X3 \in X1))) \quad (9)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(\forall X2.(m2_finseq_1 X2 X0)\Rightarrow(\forall X3.(m2_finseq_1 X3 X0)\Rightarrow((X1 \in k10_xtuple_0 X2)\Rightarrow(k1_finseq_5 X0 (k8_finseq_1 X0 X2 X3) X1 = k1_finseq_5 X0 X2 X1))))))$$