

t66_mesfunc6

(TMTA1NeYYTq4nm529toxrRp732yDGubp6WG)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_rfunct_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k19_rfunct_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ \forall X2.r2_relset_1 X0 k1_numbers (k2_partfun1 X0 k1_numbers \\ (k19_rfunct_3 X0 X1) X2) (k19_rfunct_3 X0 (k2_partfun1 X0 k1_numbers \\ X1 X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ \forall X2.r2_relset_1 X0 k1_numbers (k2_partfun1 X0 k1_numbers \\ (k18_rfunct_3 X0 X1) X2) (k18_rfunct_3 X0 (k2_partfun1 X0 k1_numbers \\ X1 X2)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.\forall X3.((m1_subset_1 X2 \\ (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1)))) \Rightarrow ((r2_relset_1 X0 X1 X2 X3) \Leftrightarrow (X2 = X3)) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2) \wedge \\ (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow ((v1_funct_1 \\ (k2_partfun1 X0 X1 X2 X3)) \wedge (m1_subset_1 (k2_partfun1 X0 X1 X2 X3) \\ (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 X1) \wedge \\ & m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow \\ & ((v1_funct_1 (k19_rfunct_3 X0 X1)) \wedge (m1_subset_1 (k19_rfunct_3 \\ & X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 X1) \wedge \\ & m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow \\ & ((v1_funct_1 (k18_rfunct_3 X0 X1)) \wedge (m1_subset_1 (k18_rfunct_3 \\ & X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \end{aligned} \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. \forall X2. ((v1_funct_1 \\ & X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow \\ & ((r2_reset_1 X0 k1_numbers (k18_rfunct_3 X0 (k2_partfun1 X0 k1_numbers \\ & X2 X1)) (k2_partfun1 X0 k1_numbers (k18_rfunct_3 X0 X2) X1)) \wedge (r2_reset_1 \\ & X0 k1_numbers (k19_rfunct_3 X0 (k2_partfun1 X0 k1_numbers X2 X1)) \\ & (k2_partfun1 X0 k1_numbers (k19_rfunct_3 X0 X2) X1)))) \end{aligned}$$